

Branching-ratio and CP violation in semi-inclusive flavor-changing top decays

L. T. Handoko*

Laboratory for Theoretical Physics and Mathematics
Indonesian Institute of Sciences

Kom. PUSPIPTEK Serpong P3FT-LIPI, Tangerang 15310, Indonesia

Abstract

Semi-inclusive top decays, $t \rightarrow q_i X_{jj}$, are examined in the framework of the standard model, with $q_i = u$ or c , X_{jj} : vector meson formed by $q_j \bar{q}_j$ ($q_j = d, s, b$). By using a simple model of hadronization, that is $\langle 0 | \bar{q}_j \gamma^\mu q_j | X_{jj} \rangle = m_{X_{jj}} f_{X_{jj}} \epsilon_{X_{jj}}^\mu$, the total branching ratio is expected to be $\mathcal{B}(t \rightarrow q_i X_{jj}) < \left(|V_{ij}^* V_{tj}| / |V_{tb}| \right)^2 \times O(10^{-3})$ depends on the size of the coupling constant $f_{X_{jj}}$. On the other hand, the CP asymmetry should be induced only by the long-distance effects of intermediate state $q_k \bar{q}_k$ ($q_k = d, s, b$) that subsequent scatters into the vector meson. The analytical results including the continuum contributions are given with keeping all masses. However, it yields the CP asymmetry to be small, i.e. $\mathcal{A}_{\text{CP}}(t \rightarrow q_i X_{jj}) < \text{Im} \left[\left(V_{ij}^* V_{tj} \right)^* (V_{ik}^* V_{tk}) \right] / |V_{ij}^* V_{tj}|^2 \times O(10^{-2})$ for an identified $q_k \bar{q}_k$ intermediate state, and suppressed more again by a factor of $\sim (m_b^2 - m_s^2) / m_{X_{jj}}^2$ for summing up all of them. Particularly, $\mathcal{A}_{\text{CP}}(t \rightarrow u \Upsilon) \simeq \pm 1\%$, induced by the intermediate state $d\bar{d}$ or $s\bar{s}$.

PACS number(s) : 14.65.Ha

*E-mail address : handoko@p3ft.lipi.go.id

After the discovery of the top quark and the determination of its mass [1], more valuable information should be expected from some classes of both tree-level and rare top decays. Because of its large mass scale, top quark will decay before hadronization and also the effects of QCD should be unambiguous and presumably very small. Then the precise determination of the modes may provide important data with less uncertainties.

In general, the rare decays are interesting because of its sensitiveness to the new physics. However, especially in the case of rare top decays, the branching ratio in the framework of standard model (SM) is largely suppressed by the GIM mechanism, e.g. $\mathcal{B}(t \rightarrow \gamma, H, g) \sim 10^{-7} \sim 10^{-10}$ as pointed out in [2]. Further, the class of these rare top decays are expected to be enhanced significantly by some new contributions due to physics beyond the SM, e.g. [2, 3]. However, they are still anticipated to be small and only in few cases [4] it is expected to be measureable at future top factories like the upgraded Tevatron and LHC [5].

The purpose of this paper is to study the semi-inclusive top decays, that is $t \rightarrow q_i X_{jj}$ with $q_i = u, c$ and X_{jj} is any vector meson formed by $q_j \bar{q}_j$ bound state as Υ for $b\bar{b}$, ϕ for $s\bar{s}$ and so on. Since the processes occur in tree-level, the measurements can be achieved easier in future experiments. The interest is, especially, focused on the CP violation in the modes by considering CP asymmetry

$$\mathcal{A}_{\text{CP}}(t \rightarrow q_i X_{jj}) \equiv \frac{(\Gamma - \bar{\Gamma})(t \rightarrow q_i X_{jj})}{(\Gamma + \bar{\Gamma})(t \rightarrow q_i X_{jj})} \equiv \frac{\Delta}{\Sigma}, \quad (1)$$

with $\bar{\Gamma}$ is the complex conjugate of Γ , while in general

$$\Delta = -2 \text{Im}(\alpha^* \beta) \text{Im}(\mathcal{M}_1^* \mathcal{M}_2), \quad (2)$$

$$\Sigma = |\alpha|^2 |\mathcal{M}_1|^2 + |\beta|^2 |\mathcal{M}_2|^2 + 2 \text{Re}(\alpha^* \beta) \text{Re}(\mathcal{M}_1^* \mathcal{M}_2), \quad (3)$$

if one writes the amplitude as $\mathcal{M} \equiv \alpha \mathcal{M}_1 + \beta \mathcal{M}_2$. Hence, the imaginary parts of $(\alpha^* \beta)$ and $(\mathcal{M}_1^* \mathcal{M}_2)$ are required to be non-zero coincidently in order to have non-zero CP asymmetry.

In the framework of the SM, the decay is realized by a tree W -exchange diagram. Then the lowest order decay amplitude of $t \rightarrow q_i X_{jj}$ is obtained,

$$\mathcal{M}^{(0)}(t \rightarrow q_i X_{jj}) = \sqrt{2} G_F (V_{ij}^* V_{tj}) \frac{f_{X_{jj}} m_{X_{jj}} m_W^2}{r^2 - m_W^2 + i \Gamma_W m_W} \epsilon_{X_{jj}}^{*\mu} [\bar{q}_i \gamma_\mu L t], \quad (4)$$

with $L = (1 - \gamma_5)/2$ and r^2 is the square of momentum transfer to W -boson. The vector meson is factorized as

$$\langle 0 | \bar{q}_j \gamma^\mu q_j | X_{jj} \rangle = m_{X_{jj}} f_{X_{jj}} \epsilon_{X_{jj}}^\mu, \quad (5)$$

where $f_{X_{jj}}$ is a constant with dimension of mass and $\epsilon_{X_{jj}}^\mu$ denotes the polarization vector. Remark that the color-factor in Eq. (4) has been omitted, and then should be considered as an unknown parameter absorbed in $f_{X_{jj}}$.

It is straightforward to obtain the total branching ratio for $t \rightarrow q_i X_{jj}$ decays by assuming the charged-current $t \rightarrow b W$ to be a dominant mode in the top quark decay. This leads

$$\mathcal{B}(t \rightarrow q_i X_{jj}) \simeq \frac{\Gamma(t \rightarrow q_i X_{jj})}{\Gamma(t \rightarrow b W)}. \quad (6)$$

In the top quark center-of-mass system, the denominator reads

$$\Gamma(t \rightarrow b W) = \frac{\hat{G}_F m_t}{8\sqrt{2}\pi} |V_{tb}|^2 \sqrt{g^{bW}} \mathcal{F}_1^{bW}, \quad (7)$$

where

$$g^{xy} \equiv 1 + \hat{m}_x^4 + \hat{m}_y^4 - 2(\hat{m}_x^2 + \hat{m}_y^2 + \hat{m}_x^2 \hat{m}_y^2), \quad (8)$$

$$\mathcal{F}_1^{xy} \equiv (1 - \hat{m}_x^2)^2 + (1 + \hat{m}_x^2) \hat{m}_y^2 - 2\hat{m}_y^4. \quad (9)$$

A caret means normalization with m_t . This agrees with some literatures in $\hat{m}_b \simeq 0$ limit. The value is, $\Gamma(t \rightarrow b W) = (1.72 \times |V_{tb}|^2)(\text{GeV})$ for $m_t = 180(\text{GeV})$, $m_W = 80.33(\text{GeV})$ and $m_b = 4.3(\text{GeV})$ [6]. Note that including the bottom-mass reduces the width by $\sim 2\%$. On the other hand, the nominator is

$$\Gamma^{(0)}(t \rightarrow q_i X_{jj}) = \frac{\hat{G}_F^2 \hat{f}_{X_{jj}}^2 \hat{m}_W^4 m_t}{8\pi} |V_{ij}^* V_{tj}|^2 \sqrt{g^{iX_{jj}}} \mathcal{F}_1^{iX_{jj}} |\mathcal{F}_2^{iX_{jj}W}|^2, \quad (10)$$

under assumptions that $E_{X_{jj}} = 2E_j$ and $m_{X_{jj}} = 2m_j$, with E denotes the time-component of four-momentum. Here $\mathcal{F}_2^{iX_{jj}W}$ is the W -boson propagator contribution,

$$\mathcal{F}_2^{xyz} \equiv \left[\frac{1}{2} (1 + \hat{m}_x^2 - \hat{m}_y^2) - \hat{m}_z^2 + i \hat{\Gamma}_z \hat{m}_z \right]^{-1}. \quad (11)$$

Then, the branching ratio for $t \rightarrow q_i X_{jj}$ decay defined in Eq. (6) is found to be, $\mathcal{B}(t \rightarrow q_i X_{jj}) \simeq 0.26 \times \hat{f}_{X_{jj}}^2 (|V_{ij}^* V_{tj}|/|V_{tb}|)^2$. Remark that the dependences on the description of the flavors (i, j) are tiny ($O(10^{-3})$ level). This means that the differences between them are mostly determined by the CKM matrix elements. Assuming $\hat{f}_{X_{jj}} \sim O(\hat{m}_{X_{jj}})$, the branching ratio can reach $\sim (|V_{ij}^* V_{tj}|/|V_{tb}|)^2 \times O(10^{-3})$.

There is no CP violation in the modes as long as considering only the lowest order diagram, because there is only one amplitude with single phase that governs the processes. So, one needs other sources with different phases to induce non-zero CP asymmetry. As mentioned before, unfortunately the loop contributions like the penguin diagrams, are also vain because of large suppression due to GIM mechanism. Then, one must proceed to consider the long-distance contributions arise from any on-mass-shell intermediate states that then scatter into X_{jj} through 1-photon exchange. Anyway, 1-gluon mediated scattering may be dominate, but unfortunately the QCD scattering can occur at the order of α_s^3 when, as the present paper, one considers only color singlet vector mesons as the final states. Of course, it may still be comparable with the electromagnetic scattering as occurred in the $q\bar{q} \rightarrow$

J/ψ [7], however since both contributions are similar and the electromagnetic one is simpler to calculate, only 1-photon mediated scattering are going to be discussed, i.e.

$$t \longrightarrow q_i (q_k \bar{q}_k) \xrightarrow{\text{QED}} X_{jj} . \quad (12)$$

Here, all intermediate states ($d\bar{d}, s\bar{s}, b\bar{b}$) can contribute to the processes. Since the top quark is not expected to form any bound states and q_i appears as a jet, the effects of the rescattering of final states, i.e. the intermediate states that have same quark content as the final states, to CP asymmetry are not important. In this meaning, the situation is different with the B decays [8, 9]. However, there will be two cases that make the size of CP asymmetries to be largely different. The first case is when one can identify and then pick up one of the intermediate states, and the second is when one can not identify and sum up all of them. In the former case the CP asymmetry can be large, while in the later one it will be suppressed by GIM mechanism. Both cases will be discussed in detail later. Further, related with the present interest, it is sufficient enough to calculate the continuum contributions of these intermediate states.

Since the long-distance contributions must be smaller by a factor of α , then it can be treated perturbatively. Hence the amplitude including one of the intermediate states can be expressed as

$$\mathcal{M}(t \rightarrow q_i X_{jj}) = \mathcal{M}^{(0)}(t \rightarrow q_i X_{jj}) + \frac{i}{2} \mathcal{M}^{(0)}(t \rightarrow q_i (q_k \bar{q}_k)) \mathcal{T}((q_k \bar{q}_k) \rightarrow X_{jj}) . \quad (13)$$

Similar with Eq. (4), $\mathcal{M}^{(0)}(t \rightarrow q_i (q_k \bar{q}_k))$ is given as

$$\mathcal{M}^{(0)}(t \rightarrow q_i (q_k \bar{q}_k)) = \sqrt{2} G_F (V_{ik}^* V_{tk}) \frac{m_W^2}{r^2 - m_W^2 + i \Gamma_W m_W} [\bar{q}_k \gamma^\mu q_k] [\bar{q}_i \gamma_\mu L t] , \quad (14)$$

and the 1-photon mediated transition amplitude is

$$\mathcal{T}((q_k \bar{q}_k) \rightarrow X_{jj}) = \alpha \pi Q_j Q_k \frac{f_{X_{jj}}}{m_{X_{jj}}} \epsilon_{X_{jj}}^\mu [\bar{q}_k \gamma_\mu q_k] . \quad (15)$$

It is obvious that the additional term in Eq. (13) induces different phase, except for $q_k = q_j$. Therefore the interference term between them will induce CP asymmetry as expected.

First, let us give the expression for the denominator in Eq. (1). Since the perturbative term in Eq. (13) is supposed to be small compared with $\mathcal{M}^{(0)}$, it can be ignored. Then the result is straightforward,

$$\Sigma \simeq \Gamma^{(0)}(t \rightarrow q_i X_{jj}) . \quad (16)$$

After that, one can consider the nominator with one intermediate state of $q_k \bar{q}_k$ corresponds with the first case mentioned above. It is calculated to be

$$\begin{aligned} \Delta_k = & - \frac{Q_j Q_k \alpha \hat{G}_F^2 \hat{f}_{X_{jj}}^2 \hat{m}_W^4 m_t}{\hat{m}_{X_{jj}}^2} \sqrt{g^{iX_{jj}}} \left| \mathcal{F}_2^{iX_{jj}W} \right|^2 \\ & \times \text{Im} \left[\left(V_{ij}^* V_{tj} \right)^* (V_{ik}^* V_{tk}) \right] \left[\mathcal{F}_3^{iX_{jj}} + \hat{m}_k^2 \mathcal{F}_4^{iX_{jj}} \right] , \end{aligned} \quad (17)$$

where the dependence of the intermediate state mass, m_k , has been collected in $\mathcal{F}_4$, that is

$$\mathcal{F}_3^{xy} = \hat{m}_y^2 \left[\frac{1}{4} (g^{xy} + 4\hat{m}_y^2) \sqrt{g^{xy} + 4\hat{m}_x^2} - (1 - \hat{m}_x^2) \sqrt{g^{xy} + 4\hat{m}_y^2} - \frac{1}{2} (1 - \hat{m}_x^2)^2 + \frac{1}{2} \hat{m}_y^4 \right], \quad (18)$$

$$\mathcal{F}_4^{xy} = 2 \left[(1 - \hat{m}_x^2 - \hat{m}_y^2) \sqrt{g^{xy} + 4\hat{m}_y^2} - (1 - \hat{m}_x^2)^2 + 2\hat{m}_y^2 \left(1 + \hat{m}_x^2 - \frac{1}{2} \hat{m}_y \right) \right]. \quad (19)$$

On the other hand, for the second case, using the unitarity relation of the CKM matrix, i.e. $\sum_{k=d,s,b} V_{ik}^* V_{tk} = 0$, and summing up all intermediate states, one finds

$$\begin{aligned} \Delta_{\text{tot}} &= \frac{1}{3} \frac{Q_j \alpha \hat{G}_F^2 \hat{f}_{X_{jj}}^2 \hat{m}_W^4 m_t}{\hat{m}_{X_{jj}}^2} \sqrt{g^{iX_{jj}}} (\mathcal{F}_2^{iX_{jj}W})^2 \mathcal{F}_4^{iX_{jj}} \\ &\times \left\{ \text{Im} \left[(V_{ij}^* V_{tj})^* (V_{ib}^* V_{tb}) \right] [\hat{m}_b^2 - \hat{m}_s^2] \right. \\ &\quad \left. - \text{Im} \left[(V_{ij}^* V_{tj})^* (V_{id}^* V_{td}) \right] [\hat{m}_s^2 - \hat{m}_d^2] \right\}, \end{aligned} \quad (20)$$

since $Q_d = Q_s = Q_b = -1/3$ and the interference terms between higher-order terms are neglected. Now, combining the results in Eqs. (16)~(20), the CP asymmetry defined in Eq. (1) can be found to be,

$$\mathcal{A}_{\text{CP}}(t \rightarrow q_i X_{jj}) \sim \begin{cases} -\mathcal{K}^{ijk} \times 0.007 & , \text{ with one } (q_k \bar{q}_k) \\ -\mathcal{K}^{ijb} \times O(10^{-5}) & , \text{ with all } (q_k \bar{q}_k) \end{cases} \quad (21)$$

where $\mathcal{K}^{ijk} \equiv \text{Im} \left[(V_{ij}^* V_{tj})^* (V_{ik}^* V_{tk}) \right] / |V_{ij}^* V_{tj}|^2$. Further, the results in Eqs. (10), (19) and (20) show that in case of summing up all intermediate states, the CP asymmetry will be suppressed by a factor of $\sim (m_b^2 - m_s^2)/m_{X_{jj}}^2$, if we put $(m_s^2 - m_d^2)/m_{X_{jj}}^2 \sim 0$. For example, definite expressions for $\mathcal{K}^{ibk}$ and $\mathcal{K}^{isk}$, corresponds with the final states Υ and ϕ , can be written as

$$\begin{aligned} \mathcal{K}^{ubd} &= \frac{-\eta}{\rho^2 + \eta^2} & , & \quad \mathcal{K}^{ubs} = \frac{\eta}{\rho^2 + \eta^2} & , & \quad \mathcal{K}^{ubb} = 0 & , \\ \mathcal{K}^{cbd} &= \lambda^2 \eta & , & \quad \mathcal{K}^{cbs} = 0 & , & \quad \mathcal{K}^{cbb} = 0 & , \\ \mathcal{K}^{usd} &= \eta & , & \quad \mathcal{K}^{uss} = 0 & , & \quad \mathcal{K}^{usb} = -\eta & , \\ \mathcal{K}^{csd} &= -\lambda^2 \eta & , & \quad \mathcal{K}^{css} = 0 & , & \quad \mathcal{K}^{csb} = 0 & , \end{aligned} \quad (22)$$

by using the Wolfenstein parametrization. The largest factor is one of the decay $t \rightarrow u (d\bar{d}, s\bar{s}) \rightarrow \Upsilon$, that is $\mathcal{K}^{ubd} = 1.65$ and $\mathcal{K}^{ubs} = -1.65$ for $(\rho, \eta) = (0.3, 0.34)$. This leads the CP asymmetry to be $|\mathcal{A}_{\text{CP}}(t \rightarrow u (d\bar{d}, s\bar{s}) \rightarrow \Upsilon)| \simeq 1\%$.

In conclusion, in the framework of the SM, it is shown that the rate of $t \rightarrow q_i X_{jj}$ may be large enough to be measured in the future top factories like LHC and the upgraded Tevatron, while the CP asymmetries are too small to be detected. However, inversely it means that the modes are clean and good probes to see CP asymmetries raise in new physics beyond the SM, for example, in the supersymmetric

(SUSY) model. The enhancement of CP asymmetry in $t \rightarrow q_i X_{jj}$ due to the contributions of scalar fermions in the SUSY model will be published in next paper [10].

The author thanks J. Hashida for useful discussion during the work. A part of this work is supported by Ministry of Education and Culture of Japan under Monbusho Fellowship Program.

References

- [1] F. Abe et.al. (CDF Collaboration), *Phys. Rev. Lett.* **74** (1995) 2626;
S. Abachi et.al. (D0 Collaboration), *ibid.* **74** (1995) 2632.
- [2] G. Eilam, J. L. Hewett and A. Soni, *Phys. Rev.* **D44** (1991) 1473.
- [3] J. L. Lopez, D. V. Nanopoulos and R. Rangarajan, *Phys. Rev.* **D56** (1997) 3100;
G. M. de Divitiis, R. Petronzio and L. Silvestrini, *Nucl. Phys.* **B504** ((1997) 45.
- [4] J. M. Yang, B-L. Young and X. Zhang, *Phys. Rev.* **D58** (1998) 055001.
- [5] *Future Electroweak Physics at the Fermilab Tevatron*, report of the TeV2000 working group, *FERMILAB-PUB 96/082* (1996) , edited by D. Amidei and R. Brock.
- [6] Review of Particle Physics, *Phys. Rev.* **D54** (1996) 1.
- [7] L. Kopke and N. Wermes, *Phys. Rep.* **174** (1989) 67 and references therein.
- [8] L. Wolfenstein, *Phys. Rev.* **D43** (1991) 151.
- [9] I. Dunietz and J. M. Soares, *Phys. Rev.* **D49** (1994) 5904;
J. M. Soares, *ibid.* **D52** (1995) 242.
- [10] L. T. Handoko and J. Hashida, *Phys. Rev.* **D58** (1998) 094008.